



Coopers Ultrakill® Wettable Powder

REG. NO. L 10064 Act No. 36 of 1947
Namibian Reg No. N-AR 1721

REG. NR. L 10064 Wet Nr. 36 van 1947
Namibiese Reg Nr. N-AR 1721

A long lasting wettable powder insecticide for the control of flying and crawling insect pests as listed.

'n Benatbare poeier insekdoder met 'n lang nawerkende aksie vir die beheer van vlieende en kruipende insekplae soos gelys.

ACTIVE INGREDIENT: / AKTIEWE BESTANDDEEL:

Alpha-cypermethrin (Pyrethroid)

50 g/kg

Alfa-sipermetrien (Piretroïed)

NET CONTENTS: / NETTO INHOUD: Contact supplier for available packaging sizes/ Kontak verskaffer vir beskikbare verpakking groottes.

REGISTRATION HOLDER/REGISTRASIEHOUER:

REGISTRATION HOLDER

Starplex 83 CC
P.O. Box 14374
BREDELL 1623
SOUTH AFRICA
+27119794246/7

Distribution by:
Coopers Environmental Science (Pty) Ltd
Reg. No. 2000/02366/70/7
91 Maple Street
POMONA, 1619
SOUTH AFRICA
TEL: +2711 979-4246/7
UN NO. 3077



CAUTION
VERSIGTIG



WARNINGS

- Handle with care.
- Store in a cool place away from food and feedstuffs.
- Keep out of reach of children, uninformed persons and animals.
- Do not apply directly onto animals.
- Do not apply to surfaces where food is handled.
- Causes severe skin and mild eye irritation.
- Poisonous when swallowed and moderately toxic by skin contact.
- Not significantly hazardous to bees, various other beneficial pest parasites and predators and fish under field conditions, provided the product is used as directed.
- Moderately toxic to wildlife.
- Do not spray over or allow drift to contaminate drinking water and water bodies such as dams, ponds, rivers, streams or fish hatcheries.

PRECAUTIONS

- Recommended protective clothing: Overall, gloves and facemask.
- Avoid skin contact and inhalation of spray mist or eye splashes.
- Wash thoroughly with soap and water after use or after accidental skin contact.
- Do not eat, drink or smoke while mixing or spraying or before washing hands and face after use.
- Wash contaminated clothing after each application.
- Contamination of food, feedstuffs, eating utensils and drinking water should be avoided.
- Remove pets and aquaria.
- Thoroughly wash and rinse the spray equipment / containers / basins after use and dispose of wash water where it will not contaminate crops, grazing, rivers and dams.
- Destroy empty container and do not re-use for any other purpose.

Although this remedy has been extensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal climatic and storage conditions, quality of dilution water, compatibility with other substances not indicated on the label and the occurrence of resistance of a pest against the remedy concerned as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to the environment or harm to man or animal or lack of performance of the remedy concerned due to failure of the user to follow the label instructions or due to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

Coopers Ultrakill® Wettable Powder spray mixture is colourless and odourless and can be used indoors or outdoors, in and around homes, factories, hotels, shops, food handling and processing establishments, hospitals, schools, dairy parlours, piggeries, stables, poultry houses, dog kennels, refuse dumps and similar places where high levels of hygiene are required.

Mixing instructions for spraying:

Half fill the sprayer with clean water. Measure the required amount of **Coopers Ultrakill® Wettable Powder** and pre-mix this with some water. Add this mixture to the water in the sprayer. Fill the sprayer with water to the required level and shake the sprayer well to ensure proper mixing or ensure continual mechanical or hydraulic agitation before spraying commences. Prepared spray mixture must not be left in the spray tank for any length of time e.g. overnight.

General directions:

Coopers Ultrakill® Wettable Powder is a long lasting wettable powder contact and stomach insecticide for the control of malaria vectors, flies, cockroaches, bedbugs, fleas, fishmoths, ants, hide beetles and -larvae as well as litter beetle and -larvae. The length of control depends on:

- insect species being treated,
- level of infestation,
- type of surface (or bed-net fabric),
- whether the treatment is applied indoors or outdoors,
- exposure to sunlight and weather conditions and,
- depending on dosage rate, months of control can be expected indoors and on bed-nets, and several weeks outdoors under favourable conditions.

Poultry uses:

- **Poultry sheds:** All cracks, crevices and potential hiding places where the litter beetle may seek shelter must receive special attention. Ensure thorough wetting of all surfaces including vertical surfaces up to at least one metre above the ground. **Coopers Ultrakill® Wettable Powder** will control but will NOT eliminate existing populations of litter beetle that have invaded and colonised the insulation of cavity walls or other protected places.
- **Broiler sheds:** Should be treated before restocking with each new batch of chicks, i.e. after the clearing, cleaning and disinfecting operations have been carried out.
- **Laying sheds:** Should be treated on a regular basis. Frequency will depend on the type of surface to be treated. Dropping and litter heaps may require treatment more often. New and empty cages can be treated BEFORE stocking with hens. DO NOT SPRAY THE BIRDS DIRECTLY OR ALLOW THE SPRAYMIST TO DRIFT OVER THEM.

MALARIA VECTOR CONTROL:

Surface sprays

- Use a sprayer fitted with a nozzle that will deliver coarse droplets at a low pressure. Do not spray at high pressure that will result in a spray mist.
- Ensure total coverage of all surfaces. All cracks, crevices and potential hiding places where mosquitoes may seek shelter must receive special attention.
- Allow spray to dry on treated surfaces before re-entering.

Bed-nets

- All nets need to be washed and dried before treatment. Some new nets are sometimes starched, and must therefore be washed first to remove the starch. Ensure that all traces of soap is rinsed out after washing and before treating with **Coopers Ultrakill® Wettable Powder**, as alkaline soap residues could influence the persistence of the treatment.
- The volume of **Coopers Ultrakill® Wettable Powder** to be used per bed-net is dependent on the length of time that control is required, as well as the size of the net.

Months of control required (dose active ingredient)	Single bed-net (±12 m ²)	Double bed-net (±16 m ²)	Queen size bed- net (±20 m ²)	King size bed-net (±24 m ²)
4 – 6 months (20 mg a.i. / m ²)	5 g	6.5 g	8 g	10 g
more than 9 months (40 mg a.i. / m ²)	10 g	13 g	16 g	20 g

- Bed-net fabric, mesh size and accessories such as a roof, border or string ties, affect the volume of water needed to treat and thoroughly wet a net. Generally 30 mℓ mixture / m² is needed for synthetic materials (e.g. polyester) and 130 – 150 mℓ / m² for cotton.
- Rough guidelines for the amount of water needed to wet different size bed-nets are given below (it is recommended however that the exact absorption volume for each type and size of net be determined before treatment).

Fabric	Single bed-net (±12 m ²)	Double bed-net (±16 m ²)	Queen size bed-net (±20 m ²)	King size bed-net (±24 m ²)
SYNTHETIC MATERIAL (e.g. Polyester)	360 mℓ	480 mℓ	600 mℓ	720 mℓ
Cotton	1700 mℓ	2250 mℓ	2800 mℓ	3400 mℓ

Mixing Instructions for treating bed-nets.

- Use a suitable container / bucket / bowl / bag, that is large enough to hold the bed-net as well as the required amount of water.
- Treat nets outside or in a well-ventilated area and use gloves.
- Fill the container with the predetermined volume of water to fully wet the bed-net to be treated. Add the correct amount of **Coopers Ultrakill® Wetable Powder** (see above) to the water and mix well.
- Soak the net in the mixture until it is thoroughly wet.
- Remove the net and allow excess to drip back into the container. (If the correct amount of mixture was used, there should be very little left).
- Allow the net to dry evenly by laying it on a bed or bedding (which helps to kill bedbugs) or on a plastic sheet outside in the shade.
- When net is dry, it may be hung up over the bed.
- After treating nets, wash hands, arms, and container and dispose of wash water where it will not contaminate crops, grazing, rivers and dams.
- The effect of the treatment will not be the same if the net is washed after treatment, and a re-treatment is advisable.
- Bed-nets treated with **Coopers Ultrakill® Wetable Powder** will not be stained or have an odour after treatment and will not be more flammable.
- To treat several identical nets, a larger volume of mixture can be prepared by multiplying the amount of **Coopers Ultrakill® Wetable Powder** as well as the volume water needed per net, by the total number of nets to be treated.

ALL SPRAY APPLICATIONS MUST BE MADE WITH SUITABLE EQUIPMENT THAT IS IN GOOD WORKING ORDER AND CORRECTLY CALIBRATED TO ENSURE THE DESIRED COVERAGE FOR THAT PARTICULAR METHOD OF APPLICATION.

PEST	DOSAGE	DIRECTIONS FOR APPLICATION
PUBLIC HEALTH: Adult house flies Adult stable flies Ants Bedbugs Cockroaches (American & German) Fishmoths Fleas	25 g (5 soluble sachets) / 5 ℓ water For high infestations – 50 g (10 soluble sachets) per 5 ℓ water	Apply as a coarse spray or with a paintbrush to cracks, crevices and any place where these insects may hide and on surfaces over which they may crawl or settle on. Apply to point of run-off. Use higher dose rate for longer lasting control or where infestation is severe. Repeat if necessary.
SKINS & HIDES Hide beetle adults and larvae	25 g (5 soluble sachets) / 5 ℓ water	Skins and Hides: Thoroughly wet all parts. Repeat treatment every 3 months or when necessary. Warehouse floors, walls and other surfaces: Apply spray mixture to the point of run-off. Repeat treatment every 3 months or if necessary.

<p>POULTRY SHEDS Lesser Mealworm / Litter beetle (<i>Alphitobius diaperinus</i>)</p>	<p>25 g (5 soluble sachets) / 5 ℓ water</p> <p>For high infestations – 50 g (10 soluble sachets) per 5 ℓ water</p>	<p>Apply as a coarse spray to the interior surfaces of the sheds. Ensure thorough wetting of the floors, walls and other surfaces. Repeat if necessary. Use higher dose rate for longer lasting control or where infestation is severe. (See under General Directions above). DO NOT SPRAY THE BIRDS DIRECTLY OR ALLOW THE SPRAYMIST TO DRIFT OVER THEM.</p>
<p>MOSQUITO ADULTS (Including malaria vectors)</p>	<p>25 g (5 soluble sachets) / 5 ℓ water (12 mg a.i. / m²)</p> <p>For a longer residual - 50 g (10 soluble sachets) per 5 ℓ water (25.5 mg a.i. / m²)</p> <p>100 g (20 soluble sachets) / 5 ℓ water (50 mg a.i. / m²)</p> <p>20 mg a.i. / m²</p> <p>OR</p> <p>40 mg / a.i. / m²</p>	<p>Apply as a coarse spray to surfaces where insects settle in areas where re-sprays can be done every 4 – 6 weeks. Apply to point of run-off.</p> <p>Use higher dose rate for longer lasting control or where infestation is severe. Repeat if necessary.</p> <p>The length of control is determined by the dosage rate and the type of treated surface. (See also GENERAL DIRECTIONS, under DIRECTIONS FOR USE above)</p> <p>MALARIA VECTOR - SURFACE TREATMENTS</p> <p>Recommended for use as a surface spray on coarse surfaces (such as clay and brick internal and external walls, roofs and ceilings, e.g. thatched). Use this rate where applications can be made only once or twice during the malaria infection period of October to May. Apply as a coarse spray to the point of run-off by using 50 mℓ spray mixture / m².</p> <p>If the applications will be made on smooth surfaces (such as glass and wood inside and outside the house) apply as a coarse spray to the point of run-off by using 25 mℓ spray mixture / m² by mixing 200 g of Coopers Ultrakill® Wetable Powder with 5 ℓ water. This will also result in applying 50 mg a.i. / m².</p> <p>Several months of control can be expected indoors at the highest dosage rates (100 g / 5 ℓ water)</p> <p>BED-NET TREATMENT</p> <p>For the required amount of Coopers Ultrakill® Wetable Powder refer to Bed-nets under MALARIA CONTROL above. See further information under GENERAL DIRECTIONS under DIRECTIONS FOR USE above as well.</p> <p>Mix the required amount of Coopers Ultrakill® Wetable Powder with enough water to wet a bed-net. Soak the net in the mixture until it is thoroughly wet. Use the higher rate for longer period of control.</p>

WAARSKUWINGS

- Hanteer versigtig.
- Bêre in 'n koel plek weg van voedsel en voer.
- Hou buite bereik van kinders, oningeligte persone en diere.
- Moenie direk op diere aanwend nie.
- Moenie aan oppervlakke waar voedsel hanteer word, aanwend nie.
- Veroorsaak erge vel- en matige oogirritasie.
- Giftig indien ingesluk en matiglik giftig deur velopname.
- Nie betekenisvol riskant vir bye, verskillende ander nuttige plaagparasiete en -predatore en vis onder veldtoestande met die verstande dat die produk gebruik word soos voorgeskryf.
- Matiglik giftig vir wild.
- Moenie oor drinkwater en watermassas soos damme, poele, riviere, strome of visteeldamme spuit, of toelaat dat spuitnewel daarheen waai nie.

VOORSORGMATREËLS

- * Voorgestelde beskermende klere: Oorpak, handskoene en gesigskerm.
- * Voorkom velkontak en inaseming van die spuitnewel, sowel as spatsels in oë.
- * Was met water en seep na gebruik of toevallige velkontak.
- * Moenie eet, drink of rook terwyl gemeng of gespuut word nie en alvorens hande en gesig nie eers gewas is nie.
- * Was gekontamineerde klere na elke aanwending.
- * Voorkom besoedeling van voedsel, voer, drinkwater en eetgerei.
- * Verwyder troeteldiere en vistenke.
- * Was en spoel spuitapparaat / houers / emmers, deeglik uit ná gebruik en doen weg met waswater op so 'n manier dat dit nie gewasse, weiding, riviere of damme sal besoedel nie.
- * Vernietig leë houers en moenie vir enige ander doel hergebruik nie.

Alhoewel hierdie middel omvattend en onder 'n groot verskeidenheid toestande getoets is, waarborg die registrasiehouer nie dat dit onder alle toestande doeltreffend sal wees nie aangesien die werking en effek daarvan beïnvloed kan word deur faktore soos abnormale klimaats- en bergingstoestande, kwaliteit van verdunningswater, saambuikbaarheid met ander stowwe wat nie op die etiket aangedui is nie, en die voorkoms van weerstand van 'n plaag teen die betrokke middel sowel as die metode, tyd en akkuraatheid van toediening. Verder aanvaar die registrasiehouer nie verantwoordelikheid vir skade aan die omgewing of vir nadelige effek op mens of dier of vir 'n gebrek aan prestasie as gevolg van die versuim van die gebruiker om etiketaanwysings na te kom of as gevolg van die ontstaan van toestande wat nie kragtens die registrasie voorsien kon word nie. Raadpleeg die verskaffer in die geval van enige onsekerheid.

GEBRUIKSAANWYSINGS: GEBRUIK SLEGS SOOS AANGEDUI

'n Spuitmengsel van **Coopers Ultrakill® Wettable Powder** is kleurloos en reukloos en kan derhalwe gespuut word buite- of binne wonings, fabriekke, hotelle, winkels, geboue waar voedsel verwerk word, hospitale, skole, melkerie, varkhokke en perdestelle, hoender- en hondhokke, vullishope en soortgelyke plekke waar 'n hoë mate van higiëne vereis word.

Menginstruksies vir bespuiting:

Maak die spuittenk halfvol met skoon water. Meet die benodigde hoeveelheid **Coopers Ultrakill® Wettable Powder** af en meng dit vooraf met 'n hoeveelheid water. Voeg hierdie mengsel by die water in die spuittenk. Vul die spuittenk met water tot by die verlangde vlak en skud die spuittenk goed om te verseker dat die spuitmengsel behoorlik gemeng is, of verseker dat aanhoudende meganiese of hidrouliese vermenging toegepas is, alvorens daar begin word met bespuiting. Aangemaakte spuitmengsel moenie in die spuittenk gelaat word vir 'n onbepaalde tyd, bv. oornag, nie.

Algemene aanwysings:

Coopers Ultrakill® Wetable Powder is 'n benatbare poeier kontak- en maaginsekdoder met 'n lang nawerkende aksie vir die beheer van malaria vektore, vlieë, kakkerlakke, weeluse, vlooië, vismotte, muskiete, velkewerlarwes en -volwassenes, miere en *Alphitobius*-kewer en -larwe. Die tydperk van beheer word beïnvloed deur;

- die insekspesie wat behandel moet word;
- die vlak van infestasië;
- die oppervlak (of bednet materiaal) wat behandel word;
- of die bespuiting binne- of buitenshuis gedoen word;
- blootstelling aan sonlig en algemene weersomstandighede en
- afhangende van die dosis, kan onder gunstige toestande, verskeie maande se beheer binnenshuis en etlike weke buitenshuis, verkry word.

Pluimvee-gebruik:

- **Pluimveehokke:** Vir effektiewe beheer van die *Alphitobius*-kewer moet veral aandag gegee word aan die kewer se moontlike wegkruipplekke soos krake en holtes. Verseker goeie benatting van sulke oppervlakke asook vertikale oppervlakke tot op 'n hoogte van ten minste een meter vanaf die grond. Indien *Alphitobius*-kewers alreeds gevestig geraak het in die mure en ander beskermende plekke, sal **Coopers Ultrakill® Wetable Powder** die getalle beheer maar NIE ten volle uitwis nie.
- **Braaikuikenhokke:** Bespuiting moet geskied nadat die hokke ontruim, skoongemaak en ontsmet is maar voordat 'n nuwe groep kuikens hervestig word.
- **Lê-hokke:** Bespuiting moet gereeld plaasvind. Die tyd tussen behandelings sal afhang van die oppervlak wat bespuit word. Mis en afvalhoë sal meer dikwels bespuit moet word. Nuwe en lê hokke kan behandel word VOORDAT hennie daarin geplaas word. SORG DAT DIE VOËLS NIE DIREK RAAKGESPUIT WORD OF DAT SPUITNEWEL NA HULLE OORWAAI NIE.

MALARIAVEKTOR BEHEER:

Oppervlakbespuitings

- Gebruik 'n spuit wat toegerus is met 'n spuitkop wat growwe druppels teen 'n lae druk kan lewer. Moet nie teen 'n te hoë druk toedien wat 'n fyn mis sal lewer nie.
- Verseker dat alle oppervlakke deeglik benat word. Alle krake, holtes en moontlike wegkruipplekke van muskiete moet veral aandag kry.
- Wag todat die bespuitte oppervlakke droog is voordat die bespuitte gebied weer binnegegaan word.

Bednette

- Alle nette moet gewas en droog wees voor behandeling. Sommige nuwe nette is gestyf, en moet ook eers gewas word. Spoel nette goed uit om van alle seep ontslae te raak, omdat seep-residue in die materiaal die na-werking van **Coopers Ultrakill® Wetable Powder** kan beïnvloed.
- Die volume **Coopers Ultrakill® Wetable Powder** wat benodig word vir die behandeling van 'n bednet is afhanklik van die lengte van beheer benodig, asook die grootte van die bednet.

Maande van beheer benodig (dosis aktiewe materiaal)	Enkel bednet ($\pm 12 \text{ m}^2$)	Dubbel bednet ($\pm 16 \text{ m}^2$)	Koningins-grootte ($\pm 20 \text{ m}^2$)	Konings-grootte ($\pm 24 \text{ m}^2$)
4 – 6 maande (20 mg a.b. / m^2)	5 g	6.5 g	8 g	10 g
meer as 9 maande (40 mg a.b. / m^2)	10 g	13 g	16 g	20 g

- Die volume water wat benodig word om 'n bednet goed te benat, is afhanklik van die grootte, die materiaal waaruit dit gemaak is en die vorm van die net. Vir sintetiese materiaal, soos poli-ester word algemeen 30 ml spuitmengsel / m^2 benodig en vir katoen tussen 130 en 150 ml / m^2 .
- In die tabel hieronder, word 'n algemene riglyn gegee van die water wat benodig word om verskillende bednette deeglik te benat (dit word egter aanbeveel om die korrekte volume water vir elke tipe bednet te bepaal voordat dit behandel word).

Materiaal	Enkel bednet (±12 m ²)	Dubbel bednet (±16 m ²)	Koningins-grootte (±20 m ²)	Konings-grootte (±24 m ²)
SINTETIES (bv. Poli-ester)	360 ml	480 ml	600 ml	720 ml
Katoen	1700 ml	2250 ml	2800 ml	3400 ml

Menginstruksies vir die behandeling van bednette.

- Gebruik 'n geskikte houër / emmer / skottel of sak, wat groot genoeg is om die bednet asook die volume water te hou.
- Behandel die nette in 'n goed geventileerde vertrek, verkieslik buitekant, en maak gebruik van handskoene.
- Vul die houër met die verlangde volume water, en voeg die nodige volume **Coopers Ultrakill® Wettable Powder** by (vir volumes benodig kyk tabelle hierbo).
- Doop die net in die mengsel totdat dit deurweek is.
- Verwyder die net en laat toe dat enige oortollige oplossing terug drup in die houër. (Indien die korrekte volume mengsel gebruik is, sal net 'n klein volume oorbly).
- Laat toe dat die net droog word deur dit plat op 'n bed of beddegoed neer te lê (op dié manier word weeluisse ook beheer). Dit kan ook buite in die skadu, op plastiek neergelê word.
- Hang dit oor die bed sodra dit droog is.
- Nadat die bednette behandel is, moet hande, arms en die houers gewas word. Doen weg met die waswater op so 'n manier dat dit nie gewasse, weiding, riviere of damme sal besoedel nie.
- Indien die net gewas word nadat dit behandel is, sal die beheer nie dieselfde wees nie en word aanbeveel dat die behandeling herhaal word.
- **Coopers Ultrakill® Wettable Powder** sal nie bednette vlek of 'n reuk nalaat nie. Nette sal ook nie meer vlambaar wees nie.
- Vir die gelyktydige behandeling van 'n aantal identiese nette, vermenigvuldig die volume water sowel as die volume **Coopers Ultrakill® Wettable Powder** wat vir een net benodig word, met die aantal nette wat behandel wil word.

Alle toedienings moet gemaak word met geskikte toerusting wat in goeie werkende toestand en korrek gekalibreer is ten einde die verlangde bedekking te gee vir die betrokke toedieningsmetode.

PLAAG	DOSIS	TOEDIENINGSVOORSKRIFTE
OPENBARE GESONDHEID Kakkerlakke (Amerikaanse en Duitse) Miere, Vismotte Vlooië Weeluisse Volwasse perdevlieë Volwasse vlieë.	25 g (5 wateroplosbare sakkies) / 5 l water Vir hoë infestاسies – 50 g (10 wateroplosbare sakkies) per 5 l water	Wend as 'n growwe bespuiting aan of dien met 'n kwas aan krake, holtes en enige plek toe waar insekte mag wegkruip of op oppervlakke waaroor insekte mag beweeg of op gaan sit. Dien tot punt van afloop toe. Gebruik die hoër dosis onder hoë infestاسiedruk of waar langer periodes van beheer tussen toedienings verlang word. Herhaal indien nodig.
VELLE EN HUIDE Velkewerlarwes Volwasse velkewers	25 g (5 wateroplosbare sakkies) / 5 l water	Velle en huide. Benat alle dele deeglik. Herhaal die behandeling elke 3 maande of indien nodig. Opbergingsperseelvoere, -mure en ander oppervlakke: Dien die spuitmengsel tot punt van afloop toe. Herhaal die behandeling elke 3 maande of indien nodig.

<p>PLUIMVEEHOKKE</p> <p>Klein meelwurm / Klein meelkewer <i>Alphitobius</i>-kewer en -larwes (<i>Alphitobius diaperinus</i>)</p>	<p>25 g (5 wateroplosbare sakkies) / 5 ℓ water</p> <p>Vir hoë infestاسies – 50 g (10 wateroplosbare sakkies) per 5 ℓ water</p>	<p>Wend aan binne-oppervlakke van hokke as 'n growwe bespuiting toe en verseker goeie benatting van vloere, mure en alle ander oppervlakke waar kewers mag voorkom. Herhaal indien nodig. Dien hoër dosis toe waar langer periodes van beheer verlang word of waar hoë infestاسie voorkom. Sien ook Algemene aanwysings hierbo. SORG DAT DIE VOËLS NIE DIREK RAAKGESPUIT WORD OF DAT SPUITNEWEL NA HULLE OORWAAI NIE.</p>
<p>VOLWASSE MUSKIETE (Malariavektore ingesluit)</p>	<p>25 g (5 wateroplosbare sakkies) / 5 ℓ water (12 mg a.b. / m²)</p> <p>Vir 'n langer nawerking – 50 g (10 wateroplosbare sakkies) per 5 ℓ water (25.5 mg a.b. / m²)</p> <p>100 g (20 wateroplosbare sakkies) / 5 ℓ water (50 mg a.b. / m²)</p> <p>20 mg a.b. / m²</p> <p>OF</p> <p>40 mg a.b. / m²</p>	<p>Die lengte van beheer word bepaal deur die dosis toegedien en die betrokke oppervlak wat behandel word. Kyk ook Algemene Aanwysings onder GEBRUIKSAANWYSINGS hierbo.</p> <p>Dien as 'n growwe bespuiting toe op oppervlakke waar insekte sit en waar bespuiting elke 4 – 6 weke herhaal kan word. Dien tot punt van afloop toe.</p> <p>Gebruik die hoër dosis onder hoë infestاسiedruk of waar langer periodes van beheer tussen toedienings verlang word. Herhaal indien nodig.</p> <p>MALARIAVEKTORE – OPPERVLAKBESPUITINGS</p> <p>Word aanbeveel as 'n oppervlakbespuiting op <u>growwe oppervlakke</u> (soos klei-, baksteen- binne en buitemure, dakke en grasdakke). Gebruik hierdie dosis waar toedienings net een of twee maal gedurende die malaria infestاسie periode van Oktober tot Mei gemaak kan word. Dien as 'n growwe bespuiting toe, deur gebruik te maak van 50 mℓ spuitmengsel / m².</p> <p>Indien die oppervlakke wat behandel sal word <u>glad</u> is, soos glas of hout, sal minder water benodig word, om die punt van afloop te bereik, (<u>25 mℓ spuitmengsel</u> / m²). In hierdie geval moet 200 g Coopers Ultrakill® Wettable Powder / 5 ℓ water gebruik word om dieselfde konsentrasie , nl. 50 mg / m² te verkry.</p> <p>Verskeie maande van beheer kan binnenshuis verwag word teen die hoër dosis (100 g / 5 ℓ water)</p> <p>BEDNETBEHANDELING</p> <p>Verwys na tabelle hierbo om die korrekte volume Coopers Ultrakill® Wettable Powder, asook volume water te bepaal, wat benodig word vir bednetbehandeling. Vir verdere inligting kyk ook onder Algemene aanwysings onder GEBRUIKSAANWYSINGS hierbo.</p> <p>Meng die benodigde volume Coopers Ultrakill® Wettable Powder met genoeg water om die bednet deeglik te benat en doop die bednet in hierdie oplossing. Gebruik die hoër dosis vir langer beheer.</p>